

A farmland biodiversity strategy is needed for China

Nationwide citizen science data show the importance of farmland outside protected areas for China's avifauna. We urge the government of China to develop a national strategy for policy and research to protect biodiversity and traditional knowledge of sustainable agriculture to meet the post-2020 goal of the Convention on Biological Diversity.

Li Li, Ruocheng Hu, Jikun Huang, Matthias Bürgi, Ziyun Zhu, Jia Zhong and Zhi Lü

Agricultural lands take up to a third of the Earth's terrestrial surface. In recent decades, the expansion and intensification of agricultural land, triggered by the continuous increase in human population and dietary changes, has become one of the direct drivers of Anthropocene biodiversity loss¹. The expansion of agricultural production, however, has neither always nor everywhere taken place at the expense of biodiversity. Instead, over the course of centuries of development of agricultural practices, a significant number of wild species have adapted or even become dependent on farmland habitats. The exact characteristics of agricultural development are, therefore, relevant to the fate of global biodiversity, and wildlife-friendly farmlands should be considered as valuable ecosystems. Half of the globe's agricultural lands are distributed in the 17 megadiverse countries, of which China possesses the largest share of the world's agricultural lands at 9.9% of the global total^{2,3}. With its long cultivation history, China has accumulated rich knowledge with regard to sustainable intensive agricultural practices at small scales. The modernization of agriculture in China in recent decades, despite its remarkable success at feeding its large population, has led to the rapid extinction of diverse land-use practices, threatening the coupling of social and ecological systems⁴. Unfortunately, China's National Biodiversity Strategies and Action Plan (NBSAP) for the Convention on Biological Diversity (CBD) fails to recognize traditional agriculture as an asset for biodiversity conservation bearing its unique cultural heritage⁵, nor the drastic change in agricultural practice as one of the causes of domestic biodiversity loss.

Overlooked biodiversity value of farmlands

Typically, the value of farmland is considered with regard to food production. China's agricultural policies primarily target




Fig. 1 | Farmland bird species in China. **a**, The crested ibis (*Nipponia nippon*) depends on rice paddy fields. Benefiting from the conservation measures of maintaining traditional land use and restricting agrochemicals, its numbers have increased from less than a dozen to over 2,000 in the past 30 years. **b**, Simulated distributions of the 220 farmland bird species in China. Six avian-diverse farming regions are numbered as follows: (1) northwest Xinjiang; (2) the Bohai Rim and parts of the North China Plain; (3) the Sichuan Basin; (4) the Poyang Lake plain and the plain of Hunan and Hubei; (5) the coastal areas and plains of the Yellow Sea; and (6) the West Guangdong and East Guangxi areas. (1) and (2) mainly consist of dry farming lands; (4) and (5) are paddy fields; (3) and (6) have both. Credit: Blickwinkel / Alamy Stock Photo (a).

this production function of farmland. In China's national implementation reports for the CBD, biodiversity in the agricultural sector related to farmland is exclusively interpreted in terms of genetic resource conservation, invasive species and pest control. The term 'biodiversity' was not mentioned in the annual budget of the Ministry of Agriculture and Rural Affairs⁶. China's newly developed land-use planning scheme, 'Major Function-Oriented Zones', aimed at optimizing the state's spatial development pattern in a top-down approach by assigning the territorial space to four types based on development intensity: development-optimized areas, development-prioritized areas, development-restricted areas and development-prohibited areas⁷.

Each area is assigned to one of the four major functions serving for industrial or urban development, food production, or ecosystem services provision⁷. Although biodiversity in development-restricted areas may benefit from strictly regulated land-use intensity, this land-use planning policy has led to the segregation of lands for nature conservation and agricultural production. In other words, under the current land-use policy arrangement in China, only lands designated as Nature Reserves are recognized for sustaining biodiversity⁸.

High biodiversity in China's farmlands

Agriculture takes the most fertile lands in China which also provide abundant resources for associated wild species (Fig.

1a). According to remote sensing data, cultivated landscapes covered 1.79 million km² in 2015⁹, that is, almost twice as much as China's national nature reserves, stretching over 0.97 million km² (ref. ¹⁰). Data to assess biodiversity distribution in China's farmlands are very limited overall. However, blooming citizen science approaches on bird species resulted in the most comprehensive nationwide avian database with fine resolution and up-to-date information on species occurrences collectively compiled by over 7,000 bird watchers¹¹. Although abundance data remain insufficient in the citizen science database, regional study in Europe has reported congruent trends of avian richness and abundance influenced by different farmland management regimes¹². Using these bird data as the best available biodiversity indicator, we simulated potential habitats of 1,111 avian species, including 167 national-protected species and 70 threatened species (see Supplementary Methods). Nearly 25% of the national-protected species and 20% of the threatened species use farmland as habitat (see Supplementary Methods and Supplementary Table 1). Looking at the top 17% (Aichi Target 11) and 50% (the 'Half-Earth' Advocacy¹²) simulated avian-species-rich pixels (1-km grain size), the ratio of cultivated landscapes or national nature reserves to the 17% and 50% pixel areas increases from 15.2% and 3.6% to 28.7% and 3.9%, respectively. The importance of the cultivated landscape is more pronounced for threatened birds, with the ratio reaching 37.6%/3.1% and 30.5%/3.6%, respectively. Among the 220 birds that use farmland for feeding or nesting, more than half have over 50% simulated potential habitats in cultivated landscapes (see Supplementary Methods and Supplementary Fig. 1). The potential habitats of the 220 species identify six avian-diverse farming regions of China: (1) northwest Xinjiang; (2) the Bohai Rim and parts of the North China Plain; (3) the Sichuan Basin; (4) the Poyang Lake plain and the plain of Hunan and Hubei; (5) the coastal areas and plains of the Yellow Sea; and (6) the West Guangdong and East Guangxi areas (Fig. 1b). Nevertheless, to date there has been a lack of both regulatory and institutional arrangements to support the enormous avian conservation value in these identified species-rich farmland regions.

CBD post-2020 framework as an opportunity for mainstreaming

China's obligation to conserve biodiversity is bound to its CBD ratification. However, among the eight megadiverse countries that have over 40% agricultural territory,

China is the only nation that lacks explicit farmland biodiversity targets in its NBSAP⁵. The only other exception is the United States, which is not a CBD signatory. Heading towards the post-2020 biodiversity framework, a broad consensus has been reached for setting conservation targets high in order to reverse the steep species-decline trajectory^{13,14}. The current global protected area system is unlikely to meet such a requirement¹⁵. In this context, the post-2020 framework may serve as an opportunity for China to review its biodiversity strategies on farmland. On the one hand, China inherits a myriad of sustainable agricultural systems that facilitate the coexistence of humans and nature¹⁶. The value of this long and rich bio-cultural heritage is a great asset for the future. On the other hand, during the same time period set for achieving the Aichi Targets, China has become the world's largest consumer of chemical fertilizer and pesticide¹⁷. The rapid agricultural intensification with heavy use of pesticides has profoundly undermined ecosystem services sustained by traditional farming¹⁸, putting both biodiversity and food safety at risk. Establishing China's farmland biodiversity strategy requires the recognition of the environmental function of agricultural land at the decision-making level. Farmland has to be valued not only for agricultural production, but also as a shared space within which a great number of species complete their life cycles, especially in areas where farmlands are small and scattered. In this sense, farmland biodiversity is one form of public good produced by agriculture as an environmental externality, which should be integrated into China's ecological compensation schemes.

To ensure domestic food security, China has set up a redline of 1.2 million km² of arable land which cannot be transformed into other types of land use. Farmland to be spared for the conservation of birds and other wild fauna and flora is rather limited at present. Thus, new agricultural systems in China should be developed to integrate both traditional wisdom and scientific knowledge of sustainable intensification towards the conservation of focal species in high-priority areas. First, a national baseline of biodiversity distributions in farmlands should be established to set spatial conservation targets, such as within the six avian-diverse farming regions in China. Second, interdisciplinary research projects should be launched to investigate the integration of traditional agricultural knowledge and modern scientific knowledge to foster the successful coupling of sustainable farming with species protection. Third, policy and market incentives should

be created to reward wildlife-friendly farming or compensate for farmers' losses in production (that is, a fund for farmland eco-compensation). Lastly, farmland biodiversity monitoring should be installed to facilitate both action- and performance-based payment. We propose that conserving farmland biodiversity should become part of the jurisdiction of state-level administrative responsibility, and research funds should be directed towards finding novel solutions for human–nature coexistence.

Reporting Summary. Further information on research design is available in the Nature Research Reporting Summary linked to this article. □

Li Li¹, Ruocheng Hu^{1,2}, Jikun Huang³, Matthias Bürgi⁴, Ziyun Zhu¹, Jia Zhong⁵ and Zhi Lü^{1,2}✉

¹Center for Nature and Society, School of Life Sciences, Peking University, Beijing, China.

²Shan Shui Conservation Center, Beijing, China.

³China Center for Agricultural Policy, School of Advanced Agricultural Sciences, Peking University, Beijing, China. ⁴Research Unit Land Change Science, Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf, Switzerland.

⁵China Birdwatching Association, Kunming, China.

✉e-mail: luzhi@pku.edu.cn

Published online: 27 March 2020
<https://doi.org/10.1038/s41559-020-1161-2>

References

- Diaz, S. et al. *Summary for Policymakers of the Global Assessment Report on Biodiversity and Ecosystem Services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services* (IPBES, 2019).
- Mittermeier, R. A. *Megadiversity: Earth's Biologically Wealthiest Nations* (Agrupacion Sierra Madre, 1997).
- Statistical Databases: Land Use Total (FAO, 2019); <http://www.fao.org/faostat/en/#data/GL>.
- Liu, Y., Duan, M. & Yu, Z. *Agric. Ecosyst. Environ.* **166**, 46–54 (2013).
- Xue, D. et al. *China National Biodiversity Conservation Strategy and Action Plan (2011–2030)* (China Environmental Science Press, 2011).
- The Annual Sectoral Budget of the Ministry of Agriculture and Rural Affairs 2019* (Ministry of Agriculture and Rural Affairs, 2019); <https://go.nature.com/2wu6G3K>.
- Fan, J., Tao, A. & Ren, Q. *J. Resour. Ecol.* **1**, 289–299 (2010).
- Li, L. & Lu, Z. Multifunctionality of land and ecological compensation: thoughts on biodiversity mainstreaming. *Resour. Ind. Econ.* **7**, 12–17 (2019).
- Xu, X. et al. *China's Multi-Period Land Use Land Cover Remote Sensing Monitoring Data Set (CNLUCC)* (Resource and Environment Data Cloud Platform, 2018).
- 2018 Report on the State of the Ecology and Environment in China (Ministry of Ecology and Environment, 2019); <https://go.nature.com/3cwWkAj>.
- Hu, R. et al. *Biol. Conserv.* **211**, 47–55 (2017).
- Wilson, E. O. *Half-Earth: Our Planet's Fight for Life* (WW Norton & Company, 2016).
- Dinerstein, E. et al. *Sci. Adv.* **5**, eaaw2869 (2019).
- Mace, G. M. et al. *Nat. Sustain.* **1**, 448–451 (2018).
- Visconti, P. et al. *Science* **364**, 239–241 (2019).
- Koohafkan, P. in *Sustainable Diets: Linking Nutrition and Food Systems* (eds Burlingame, B. & Dernini, S.) Ch. 21 (CABI, 2018).
- National Bureau of Statistics of the People's Republic of China *China Statistical Yearbook 2018* (China Statistics Press, 2018).
- Huang, J. et al. *Environ. Res. Lett.* **13**, 064027 (2018).

Acknowledgements

We are grateful to F. Hua, F. Fasnacht, D. Wilcove and F. Herzog for valuable discussions and comments. G. Gautschi helped us with the English editing of the manuscript.

Author contributions

Z.L., L.L., J.H. and M.B. conceptualized and framed the main perspective of the Comment. J.Z. organized

nationwide birding records of China. R.H. conducted all species modelling and mapping and wrote the chapter of Supplementary Methods. Z.Z. reviewed the current agricultural policies of China. J.H. provided recommendations regarding agricultural policies. L.L. drafted the manuscript; M.B., Z.L. and J.H. revised the content and language of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41559-020-1161-2>.

Reporting Summary

Nature Research wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Research policies, see [Authors & Referees](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a Confirmed

- The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- A description of all covariates tested
- A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection

Data analysis

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors/reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Research [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A list of figures that have associated raw data
- A description of any restrictions on data availability

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

- Life sciences Behavioural & social sciences Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see nature.com/documents/nr-reporting-summary-flat.pdf

Ecological, evolutionary & environmental sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	To assess avian diversity distributions in China's farmland, we used MaxEnt 3.4.1 to model range maps of birds in China based on citizen science data. 160,086 independent localities of 1,111 bird species were used for the modelling.
Research sample	For each species, the MaxEnt model was performed to derive their distribution range in farmland. Taking the most widespread hawk in China's farmland for example, we collected 479 independent localities of <i>Accipiter nisus</i> , the model results showed that the habitat in farmland contains 443,413 pixels. The test AUC is 0.832, and the training AUC 0.853.
Sampling strategy	Occurrence data of 1,111 birds were collected from the citizen science project, Bird Report (www.birdreport.cn). To ensure the accuracy of location and species identification, each submitted record was checked by an experienced reviewer to filter out questionable entries. Only birdwatchers who have submitted more than 300 bird species and 100 birding reports in China are eligible to become reviewers. Collecting 47 thousand birding reports (one report may contain 1-120 species records) from 1998 to 2017, we verified the accurate coordinates of each bird species record using Google Maps API.
Data collection	All birding data from 1998 to 2017 were collected from birdreport.cn by Ruocheng Hu. Bird Report is the only large-scale nationwide project for submitting birdwatching records in China, with records of more than 1,390 bird species from 1998 to 2017. To ensure the accuracy of location and species identification, each submitted record was checked by an experienced reviewer to filter out questionable entries.
Timing and spatial scale	Original birding data were collected for the time period 1998 to 2017 from all over China.
Data exclusions	We corrected statistical errors, including over-fitting and high-variability predictions: 1) Bird species with fewer than five independent localities were excluded; 2) when localities were clustered for a species (i.e., the distance between any two records was less than one arc-minute), we randomly remove the extra locality when conducting the habitat modeling. The exclusion criteria were pre-established and were applied consistently throughout the analysis.
Reproducibility	The MaxEnt model was set by 10 cross-validated replicates for each species. Consistent results can be derived for each habitat range model.
Randomization	We modelled distribution range for all species in the citizen science database of China. The randomization was not relevant to our study except that when localities were clustered for a species (i.e., the distance between any two records was less than one arc-minute), we randomly remove the extra locality when conducting the habitat modeling.
Blinding	In the MaxEnt modeling, we randomly set 80% data for training and 20% data for testing to produce the final distribution map.
Did the study involve field work?	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

n/a	Involvement in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> Antibodies
<input checked="" type="checkbox"/>	<input type="checkbox"/> Eukaryotic cell lines
<input checked="" type="checkbox"/>	<input type="checkbox"/> Palaeontology
<input checked="" type="checkbox"/>	<input type="checkbox"/> Animals and other organisms
<input checked="" type="checkbox"/>	<input type="checkbox"/> Human research participants
<input checked="" type="checkbox"/>	<input type="checkbox"/> Clinical data

Methods

n/a	Involvement in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> ChIP-seq
<input checked="" type="checkbox"/>	<input type="checkbox"/> Flow cytometry
<input checked="" type="checkbox"/>	<input type="checkbox"/> MRI-based neuroimaging